

学校编码：10384

学号：21720071154115

廈門大學

硕士学位论文

文昌鱼GMNN基因的克隆、序列分析、表征及表达

Cloning, Sequencing, Characterization and
Expression of GMNN Gene from Amphioxus

Emmanuel MUKWAYA

指导教师：王义权

专业名称：化学生物学

答辩日期：2009年6月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1. 经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

摘 要

Geminin基因 (GMNN) 在抑制再复制的进程中是主要的调控者, 在细胞周期中的S, G2和早中期时, 它负向调节Cdt1p, 阻止Mcm2 -7p复合物在复制起点的装配, 因此GMNN通过阻止DNA非正常地再复制来维持基因组的忠实性, 确保同一个复制起点在一个细胞周期中不会活化两次。在后生动物后期生长中, GMNN在控制细胞分化的同时也控制着细胞的增殖。

GMNN分别抑制青鳉鱼视网膜中Six3和小鼠胚胎中Hox的转录因子, 同时在非洲爪蟾中通过抑制Brg-1的活性来维持神经祖细胞处于未分化状态。我们从文昌鱼*B. japonicum*中克隆得到GMNN的cDNA克隆。运用5' RACE和3' RACE克隆得到GMNN的全长cDNA 为721 bp, 经进一步测序验证, 然后运用Real time- PCR研究其在胚胎和组织中的表达情况。

文昌鱼GMNN序列编码一个241个氨基酸的蛋白质, 它与人GMNN 基因的同源性接近48.02%, 与小鼠GMNN基因的同源性达到53.73%, 与大鼠GMNN基因的同源性达到62.22%, 与原鸡GMNN基因的同源性达到52.4%, 与倭黑猩猩GMNN基因的同源性达到62.50 %。通过对文昌鱼的GMNN基因的研究我们发现它是由8个外显子和7个内含子组成, 此外, 运用实时定量PCR研究GMNN基因的表达情况显示它在卵巢中高表达, 而在精巢没有得到高表达。文昌鱼GMNN基因的克隆和表达结果显示文昌鱼该基因是一个编码相对短的蛋白的小基因, 而与其他脊椎动物不同的是该基因并不在增生组织表达。

关键词：文昌鱼；GMNN基因；Cdt1p；Mcm2 -7p复合物

Abstract

Geminin gene (GMNN) is a central regulator of the process that inhibits re-replication. During S, G2, and early M phases of the cell cycle, it negatively regulates the Cdt1p and prevents them from loading the Mcm2–7p complex to the origins of replication. GMNN thus maintains the integrity of the genome by preventing abnormal re-replication of DNA. It ensures that, the same origin of replication is not activated twice within the same cell cycle. GMNN gene controls differentiation as well as cell proliferation during late metazoan development.

It inhibits both Six3 and Hox transcription factors in Medaka retina and the mouse embryo, respectively, and maintains an undifferentiated state in neural progenitor cells of *Xenopus laevis* by antagonizing Brg-1 activity.

We have cloned a cDNA clone encoding GMNN from the amphioxus, *B. japonicum*. The 721 bp full-length GMNN gene cDNA was cloned using 5'RACE and 3' RACE and further confirmed by nucleotide sequencing and then expressed using Real time RT-PCR.

The amphioxus GMNN gene sequence encodes a protein of 241 amino acids which displays approximately 48.02% homology with human GMNN gene, 53.73% with the mouse GMNN gene, 62.22% with *Rattus norvegicus* GMNN gene, 52.4% with the *Gallus gallus* GMNN gene and 62.50 % with *Pan troglodytes*.

By studying GMNN gene sequence organization we found that it is made of 8 exons and 7 introns. Furthermore, the expression pattern of GMNN gene with Real Time RT-PCR revealed that GMNN gene is highly expressed in the ovary but not in the testis. Our results on cloning and expression the GMNN gene in amphioxus revealed that the amphioxus GMNN gene is a short gene encoding a relatively short protein but unlike in other vertebrates, is not expressed in proliferative tissues.

Keywords: Amphioxus; GMNN gene; Cdt1p; Mcm2–7p complex

厦门大学博硕士论文摘要库

参考资料

- [1] Holland, N. D. & Chen, J. Origin and early evolution of vertebrates: new insights from advances in molecular biology, anatomy, and paleontology. *News and Rev. in Mol. Cell and Dev. Biol.*, 2001, 23:142-151.
- [2] Strokes M. and Holland N. Reproduction of the Florida lancelet (*Branchiostoma floridae*): spawning patterns and fluctuations in gonad indexes and nutritional reserves. *Invertebrate Biology*, 1996, 115: 349-359.
- [3] Luan, J., et al. Characterization, Evolution and Expression of the Calmodulin1 Genes from the Amphioxus *Branchiostoma belcheri tsingtauense*. *Acta Biochimica et Biophysica Sinica*, 2007, 39:255-264.
- [4] Lacalli T.C. Prospective protochordate homologs of vertebrate midbrain and MHB, with some thoughts on MHB origins. *Int. J. Biol. Sci.*, 2006, 2: 104-9.
- [5] Minguillón C. et al. The amphioxus Hairy family: differential fate after duplication. *Development* 2003, 130: 5903-14.
- [6] Panopoulou G. et al. New evidence for genome-wide duplication at the origin of vertebrate using an amphioxus gene set and completed animal genomes. *Gen. Res.*, 2003, 13:1056-1066.
- [7] Kohn M. Reconstruction of A my-old ancestral vertebrate protokaryote. *Trends Genet.* 2006, 22:203-210.
- [8] Holland L.Z. and Yu J.K. Cephalochordate (amphioxus) embryos: procurement, culture, and basic methods. *Methods Cell Biol.* 2004, 74: 195-215.
- [9] Holland L.Z., et al. The chordate amphioxus: an emerging model organism for developmental biology. *Cell Mol Life Sci.* 2004, 61: 2290-308.
- [10] Holland and Garcia-Fernandez . Hox genes and chordate evolution. *Dev. Biol.* 1996, 173:382 – 395.
- [11] <http://www.earthlife.net/inverts/cephalochordata.html>.
- [12] Wada H. and Satoh N. Details of the evolutionary history from invertebrates to vertebrates, as deduced from the sequences of 18S rDNA. *Proc. Natl. Acad. Sci. USA*, 1994, 91:1801 – 1804.
- [13] Holland L.Z. et al. The amphioxus genome illuminates vertebrate origins and cephalochordate biology. *Genome Res.*, 2008, 18(7):1100-1111.
- [14] Holland N.D. Sequence and developmental expression of *AmphiDII*, an amphioxus *Distal-less* gene transcribed in the ectoderm, epidermis and nervous system: Insights into evolution of craniate forebrain and neural crest. *Development*, 1996, 122: 2911 – 2920.
- [15] Shimeld S.M. and Holland P.W.H. Special Feature: Vertebrate Innovations. *Proc Natl Acad Sci U S A*, 2000, 97: 4449-4452.
- [16] Meulemans D. and Bronner-Fraser M. Gene-regulatory interactions in neural crest evolution and development. *Dev. Cell*, 2004, 7: 291 – 299.
- [17] Dutta A. and Bell S.P. Initiation of DNA replication in eukaryotic cells. *Annu. Rev. Cell Dev. Biol.*, 1997, 13:293-32.
- [18] Kelly, T.J. & Brown, G.W. Regulation of chromosome replication. *Annu. Rev. Biochem.* 2000, 69:829-880.
- [19] McGarry T.J. Geminin deficiency causes a Chk1-dependent G2 arrest in *Xenopus*. *Mol. Biol. Cell*, 2002, 13: 3662-3671.
- [20] Wohlschlegel J. et al. Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Science*, 2000, 290: 2309-2312
- [21] Quinn L.M. et al. The *Drosophila* Geminin homolog: roles for Geminin in limiting DNA replication, in anaphase and in neurogenesis. *Gen. Dev.* 2001, 15: 2741-2754.
- [22] Tada S., et al. Repression of origin assembly in metaphase depends on inhibition of RLF-B/Cdt1 by geminin. *Nat. Cell Biol.*, 2001, 3:107-113.
- [23] Nyberg K.A. et al. Toward maintaining the genome: DNA damage and replication checkpoints. *Annu. Rev. Genet.* 2002, 36:617-656.
- [24] McGarry T.J. and Kirschner M.V. .Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell*, 1998, 93:1043 – 1053.

- [25] McGarry T.J. Geminin Deficiency Causes a Chk1-dependent G2 Arrest in *Xenopus*. *Mol. Biol. Cell* October 1, 2002, 13: 3662-3671.
- [26] Mihaylov I.S. et al. Control of DNA replication and chromosome ploidy by geminin and cyclin. *Mol. Cell. Biol.* 2002, 22: 1868 – 1880.
- [27] Lee C. et al. Structural basis for inhibition of the replication licensing factor: Cdt1 by GMNN. *Nature*, 2004, 430:913 – 917.
- [28] Nishitani H. et al. Proteolysis of DNA replication licensing factor Cdt1 in S-phase is performed independently of geminin through its N-terminal region. *J. Biol. Chem.* 2004, 279:30807-30816.
- [29] Donovan S. et al. Cdc6p-dependent loading of Mcm proteins onto pre-replicative chromatin in budding yeast. *Proc. Natl. Acad. Sci.* 1997, 94: 5611 – 5616.
- [30] Hofmann J. F., and Beach D. 1994. Cdt1 is an essential target of the Cdc10/Sct1 transcription factor: requirement for DNA replication and inhibition of mitosis. *Embo.J.* 1994, 13:425-434.
- [31] Gee H. Vertebrates morphology: Return of the amphioxius. *Nature*, 1994, 370: 504-505.
- [32] Rathjen et al. Pluripotent cell division cycles are driven by ectopic Cdk2, cyclin A/E and E2F activities. *Oncogene* 2002, 21: 8320 – 8333.
- [33] Weinreich et al. The Cdc6p nucleotide-binding motif is required for loading Mcm proteins onto chromatin. *Proc. Natl. Acad. Sci. U. S. A.* 1999, 96, 441 – 446.
- [34] Whittaker A.J., et al. *Drosophila* double parked: A conserved, essential replication protein that colocalizes with the origin recognition complex and links DNA replication with mitosis and the down-regulation of S phase transcripts. *Genes & Dev.* 2002, 14, 1765-1776.
- [35] Threpaut M. et al. Crystallization and preliminary X-ray crystallographic analysis of human geminin coiled-coiled domain. *Biochim. Biophys. Acta.* 2002, 1599:149-151.
- [36] Kroll K.L. et al. Geminin a neuralizing molecule that demarcates the future neural plate at the onset of gastrulation. *Dev.* 1998, 125:3247-3258.
- [37] Filippo D. B. et al. Direct interaction of geminin and Six3 in eye development. *Nature*, 2004, 427: 745 – 749.
- [38] Hara K., et al. Geminin is essential for the development of preimplantation mouse embryos. *Genes Cells*, 2006 11: 1281 – 1293.
- [39] Blow J.J. and Laskey R. A. A role for the nuclear envelope in controlling DNA replication within the cell cycle. *Nature* , 1998, 332:546 – 548.
- [40] Stillman B. Cell cycle control of DNA replication. *Science*, 1996, 274:1659 – 1664.
- [41] Tye B.K. MCM proteins in DNA replication. *Annu. Rev. Biochem.* 1999, 68, 649 – 686.
- [42] Bell S.P. & Stillman B. ATP-dependent recognition of eukaryotic origins of DNA replication by a multiprotein complex. *Nature* 1992, 6374:128 – 134.
- [43] Diffley J.F. et al. Two steps in the assembly of complexes at yeast replication origins in vivo. *Cell*, 1994, 78, 303 – 316.
- [44] Coleman T.R. et al. The *Xenopus* Cdc6 protein is essential for the initiation of a single round of DNA replication in cell-free extracts. *Cell*, 1996, 87:53 – 63. *Development*, 1996, 124, 1723 – 1732.
- [45] Gillespie PJ, Blow JJ. Nucleoplasmin-mediated chromatin remodelling is required for *Xenopus* sperm nuclei to become licensed for DNA replication. *Nucleic Acids Res.* 2000, 28: 472-480.
- [46] Maiorano D. et al. CDT1 is required for the assembly of pre-replicative complexes in *Xenopus laevis*. *Nature*, 2000, 404: 622 – 625.
- [47] Blow J.J. and Tada S. A new check on issuing the license. *Nature*, 2000, 404: 560-561.
- [48] Zou L. and Stillman B. Formation of a preinitiation complex by S-phase cyclin CDK-dependent loading of Cdc45p onto chromatin. *Science*, 1998, 280, 593 – 596.
- [49] Walter, J. & Newport, J. (2000) Initiation of eukaryotic DNA replication: origin unwinding and sequential chromatin association of Cdc45, RPA, and DNA polymerase alpha. *Mol. Cells* 5, 617 – 627.
- [50] Muzi F.M. et al. Cdc18+ regulates initiation of DNA replication in *Schizosaccharomyces pombe*. *Proc Natl. Acad. U S A.* 1996, 93: 1566 – 1570.

- [51] Drury L.S. et al. The Cdc4/34/53 pathway targets Cdc6p for proteolysis in budding yeast. *EMBO J.* 1997, 16:5966 – 5976.
- [52] Baum B., et al. Cdc18 transcription and proteolysis couple S phase to passage through mitosis. *EMBO J.* 1998, 19:5689 – 5698.
- [53] Kelly T.J. et al. The fission yeast *cdc18+* gene product couples S phase to START and mitosis. *Cell*, 1993, 74: 371 – 382.
- [54] Nishitani H. et al. The Cdt1 protein is required to license DNA for replication in fission yeast. *Nature*, 2000, 404, 625 – 628.
- [55] Lygerou Z. and Nurse P: Cell cycle. License withheld: geminin blocks DNA replication. *Science* 2000, 290:2271 – 2273.
- [56] Hendrickson et al., 2006. Phosphorylation of mcm4 by cdc2 protein kinase inhibits the activity of the minichromosome maintenance complex. *Proc. Natl. Acad. Sci. U S A*, 93(22):12223 – 12228.
- [57] Aparicio O.M., et al., 1997. Components and dynamics of DNA replication complexes in *S. cerevisiae*: Redistribution of MCM proteins and Cdc45p during S phase. *Cell*, 91: 59 – 69.
- [58] Parisi, T. et al. 2003. Cyclins E1 and E2 are required for endoreplication in placental trophoblast giant cells. *EMBO J.* 2003, 22: 4794 – 4803.
- [59] Labib K., et al. G1-phase and B-type cyclins exclude the DNA-replication factor Mcm4 from the nucleus. *Nat. Cell Biol.* 1999, 1: 415 – 422.
- [60] Nguyen V.Q. et al. Clb/Cdc28 kinases promote nuclear export of the replication initiator proteins Mcm2-7. *Curr. Biol.* 2000, 10: 195-205.
- [61] Seo S. et al. Geminin regulates neuronal differentiation by antagonizing Brg1 activity. *Genes Dev.* 2005, 19: 1723 – 1734.
- [62] Lei M., and Tye B.K. Initiating DNA synthesis: from recruiting to activating the MCM complex. *J. Cell Sci.*, 2001, 114:1447-1454.
- [63] Ralston A. and Rossant, J. Genetic regulation of stem cell origins in the mouse embryo. *Clin. Genet.* 2005, 68: 106 – 112.
- [64] Geng, Y., et al. Cyclin E ablation in the mouse. *Cell*, 2003, 114: 431 – 443.
- [65] Luo L. et al. The cell-cycle regulator geminin inhibits Hox function through direct and polycomb-mediated interactions. *Nature*, 2004, 427:749 – 753.
- [66] Kenichi Y. et al. The destruction box of human Geminin is critical for proliferation and tumor growth in human colon cancer cells. *Oncogene*, 2001, 23, 58 – 70
- [67] Kristen L.K. Geminin in embryonic development: coordinating transcription and the cell cycle during differentiation. *Frontiers in Bioscience*, 2007, 12:1395-1409
- [68] Dudderidge T. J., et al. Geminin and KI67 define proliferative state and are prognostic markers in renal cell carcinoma. *Clin Cancer Res.*, 2005, 11, 2510-7
- [69] Nilima B. et al. Human Cytomegalovirus Infection leads to Accumulation of Geminin and Inhibition of the Licensing of Cellular DNA Replication. *Journal of virology*, 2001, 77: 2369 – 2376.
- [70] Yoshida K. and Inoue I. Regulation of Geminin and Cdt1 expression by E2F transcription factors. *Oncogene*, 2004; 23:3802 – 3812
- [71] Vaziri C. et al. A p53 dependent checkpoint pathway prevents rereplication. *Mol. Cell*, 2003, 11, 997 – 1008.
- [72] Yanagi K. et al. *Caenorhabditis elegans* geminin homologue participates in cell cycle regulation and germ line development. *J. Biol. Chem.* 2005, 280: 19689 – 19694.
- [73] Michael A. et al., 2006, Geminin is essential to prevent endoreduplication and to form pluripotent cells during mammalian development, *Genes Dev.* 20:1880-1884.
- [74] Saitou N. and Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.*, 1987, 4: 406-425.
- [75] Kumar S. et al. MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics*, 2001, 17: 1244-1245.

- [76] Thompson J.D. et al. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994 22: 4673-4680.
- [77] Putnam N. H. et al., 2008. The amphioxus genome and the evolution of the chordate karyotype. *Nature* 453:1064-1072.
- [78] Ruvinsky I., et al. Phylogenetic Analysis of T-Box Genes Demonstrates the Importance of Amphioxus for Understanding Evolution of the Vertebrate Genome Genetics, 2000, 156(3): 1249-1257.
- [79] Thomas J., et al. 1998. Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell*, 1998, 93: 1043 – 1053.
- [80] [Http:// www.ncbi.nlm.nih.gov/](http://www.ncbi.nlm.nih.gov/).
- [81] [Http://genome.jgi-psf.org/Brafl1/Brafl1.home.html](http://genome.jgi-psf.org/Brafl1/Brafl1.home.html).
- [82] [Http://www.bio.davidson.edu/Courses/Molbio/MolStudents/spring2003/Pierce/realtimepcr.htm](http://www.bio.davidson.edu/Courses/Molbio/MolStudents/spring2003/Pierce/realtimepcr.htm).
- [83] James A. et al. Expression of geminin as a marker of cell proliferation in normal tissues and malignancies. *Am. Path.* 2002, 161 (1):267-274.
- [84] Xouri G. Z. et al. Cdt1 and geminin are downregulated upon cell cycle exit and are over-expressed in cancer-derived cell lines. *Eur. J. Biochem.* 2004, 271: 3368-78.
- [85] Montanari M., et al. Increased expression of geminin stimulates the growth of mammary epithelial cells and is a frequent event in human tumors. *J. Cell Physiol.* 2005, 202:215-22.

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库